

# CHAPTER 3

## LINEAR SYSTEMS & MATRICES




### SEC 3.1. INTRO TO LINEAR SYSTEMS

DEFINITION A linear equation in the variables  $x_1, x_2, \dots, x_n$  is of the form

$$a_1 x_1 + a_2 x_2 + \dots + a_n x_n = b$$

KNOWN CONSTANTS

### EXAMPLES

# VARS	VAR NAMES	EQUATION	A GEOMETRICAL INTERPRETATION OF SOLN
1	$x$	$a_1 x = b$	 Point in 1D space
2	$x, y$	$a_1 x + a_2 y = b$	 line in 2D space
3	$x, y, z$	$a_1 x + a_2 y + a_3 z = b$	 Plane in 2D space

$n$  $x_1, \dots, x_n$ 

$a_1 x_1 + \dots + a_n x_n = b$

 $(n-1)$ D plane in  $n$ D space

DEFINITION A system of linear equations in  $n$  variables is a finite collection of linear equations. Their solutions are  $n$ -tuples  $(x_1, \dots, x_n)$  that satisfy all equations simultaneously.


## EXAMPLES

(1) A system of 2 lin eqns in 2 variables, say  $x, y$ , looks like


$$\begin{cases} a_{11}x + a_{12}y = b_1 \\ a_{21}x + a_{22}y = b_2 \end{cases}$$

The solutions to each of these form a straight line in the  $xy$  plane. There are 3 different scenarios for solutions to this system:


(1) the lines are not parallel: there is 1 solution



(2) the lines are parallel & equal:  $\infty$  number of solutions



(3) the lines are parallel & not equal: no solutions



DEFINITION If a system has at least 1 solution we call it consistent and otherwise we call it inconsistent.

How to solve a system of linear equations

→ METHOD OF ELIMINATION

The method of elimination is based on the fact that the following operations don't change the solutions to a system of equations:

(1) Swapping 2 equations. This only changes the position of the eqns in the list but not their solutions

EXAMPLE  $\begin{cases} 5x + 7y = 3 \\ -x + y = 4 \end{cases}$  has same solutions as  $\begin{cases} -x + y = 4 \\ 5x + 7y = 3 \end{cases}$

(2) Multiplying an equation by a NONZERO constant. Since, later on, you can always divide by that constant again to retrieve the old equation, this doesn't change its solutions.

EXAMPLE  $\begin{cases} 5x + 7y = 3 \\ -x + y = 4 \end{cases}$  has same solutions as  $\begin{cases} 10x + 14y = 6 \\ -x + y = 4 \end{cases}$

(3) Adding a multiple of one equation to another while keeping the original equation intact. This is essentially the same as saying: you can add the same number to both sides of an equation without changing its solutions.

**EXAMPLE**  $\begin{cases} 5x + 7y = 3 \\ -x + y = 4 \end{cases}$  has same solutions as  $\begin{cases} 5x + 7y = 3 \\ 4x + 8y = 7 \end{cases} \leftarrow \text{Eqn 1} + \text{Eqn 2}$

Even though we're not adding numbers, but rather expressions involving variables that look different on both sides of the equation, (like  $5x + 7y$  and  $3$ ) we are given that these are equal ( $5x + 7y = 3$ ) and it's therefore perfectly fine to add these "different looking" terms to both sides of the equation.

Using these operations we can eliminate variables from equations in a systematic way

## EXAMPLES

(1) So we  $\begin{cases} x - 2y = 8 \quad (E1) \\ 5x + 3y = 1 \quad (E2) \end{cases}$

By subtracting 5 times the first equation from the second we get

$$\begin{cases} x - 2y = 8 \\ 5x + 3y = 1 \end{cases} \xrightarrow{(E_2) - 5(E_1)} \begin{cases} x - 2y = 8 \quad (E_1) \\ 0x + 13y = -39 \quad (E_2) \end{cases}$$

Now we can multiply both sides of (E2) by  $\frac{1}{13}$  to get

$$\begin{cases} x - 2y = 8 \\ 13y = -39 \end{cases} \xrightarrow{\frac{1}{13}(E_2)} \begin{cases} x - 2y = 8 \\ y = -3 \end{cases}$$

Now we can add (E2) twice to (E1) to get

$$\begin{cases} x - 2y = 8 \\ y = -3 \end{cases} \xrightarrow{(E_1) + 2(E_2)} \begin{cases} x = 2 \\ y = -3 \end{cases}$$

so our solution is  $(x, y) = (2, -3)$

(2) Solve  $\begin{cases} 2x + 6y = 4 \\ 3x + 9y = 11 \end{cases} \xrightarrow{(E_1) \cdot \frac{1}{2}} \begin{cases} x + 3y = 2 \\ 3x + 9y = 11 \end{cases}$

$$\xrightarrow{(E_2) - 3(E_1)} \begin{cases} x + 3y = 2 \\ 0 + 0 = 5 \quad \text{⚡} \end{cases}$$

(3) Solve  $\begin{cases} 2x + 6y = 4 \\ 3x + 9y = 6 \end{cases} \xrightarrow{(E_1) \cdot \frac{1}{2}} \begin{cases} x + 3y = 2 \\ 3x + 9y = 6 \end{cases}$

$$\xrightarrow{(E_2) - 3(E_1)} \begin{cases} x + 3y = 2 \\ 0 + 0 = 0 \end{cases} \xrightarrow{\text{always satisfied}} \{ x + 3y = 2. \leftarrow \text{for each } x \text{ there is } y \text{ that solves it}$$


In the case where we have an  $\infty$  number of solutions it is customary to describe the solutions using one (or more) parameter(s).

Let  $x = t$   $\leftarrow$  can be any real number, then  $t + 3y = 2$  so  $y = \frac{2-t}{3} = \frac{2}{3} - \frac{1}{3}t$

All solutions to the system can therefore be written

as  $(x, y) = (t, \frac{2}{3} - \frac{1}{3}t)$  or equivalently  
 complex are added piecewise

$$(x, y) = (0, \frac{2}{3}) + t \cdot \underbrace{\left(1, -\frac{1}{3}\right)}_{= (t, -\frac{1}{3}t)} \quad \begin{matrix} (a, b) + (c, d) \\ = (a+c, b+d) \end{matrix}$$



Note you could also have chosen  $y = t$ , in which case  $x = 2 - 3t$  and our solutions look like

$(x, y) = (2 - 3t, t)$ . While these look different from the solution set as a whole is the same

(4) Solve:

$$\begin{cases} x + 2y + z = 4 & E_1 \\ 3x + 8y + 7z = 20 & E_2 \\ 2x + 7y + 9z = 23 & E_3 \end{cases}$$

First we use  $E_1$  to eliminate  $x$  from  $E_2$  and  $E_3$

$$\begin{matrix} (E_2) - 3(E_1) \\ (E_3) - 2(E_1) \end{matrix} \rightarrow \begin{cases} x + 2y + z = 4 \\ 2y + 4z = 8 \\ 3y + 7z = 15 \end{cases} \xrightarrow{(E_2) \cdot \frac{1}{2}} \begin{cases} x + 2y + z = 4 \\ y + 2z = 4 \\ 3y + 7z = 15 \end{cases}$$

Now we use  $(E_2)$  to get rid of  $y$  in  $(E_3)$ . We can't use  $(E_1)$  for this since then we would introduce  $x$  again!

$$\xrightarrow{(E_3) - 3(E_2)} \begin{cases} x + 2y + z = 4 \\ y + 2z = 4 \\ z = 3 \end{cases}$$

Now we can use  $(E_3)$  to eliminate  $z$  from  $(E_1)$  and  $(E_2)$

$$\xrightarrow{\begin{matrix} (E_1) - (E_3) \\ (E_2) - 2(E_3) \end{matrix}} \begin{cases} x + 2y = 1 \\ y = -2 \\ z = 3 \end{cases} \xrightarrow{E_1 - 2(E_2)} \begin{cases} x = 5 \\ y = -2 \\ z = 3 \end{cases}$$

(5) Solve 
$$\begin{cases} 3x - 8y + 10z = 23 \\ x - 3y + 2z = 5 \\ 2x - 9y - 8z = -11 \end{cases}$$