

# SEC 8.5 INVERSES OF MATRICES

Remember: to bring matrix in REF we applied row operations, e.g.

$$\begin{array}{c} \underline{\underline{A}} \\ \begin{array}{|ccc|c} \hline 1 & 2 & 3 & 4 \\ \hline 0 & 1 & 2 & 3 \\ \hline 1 & 5 & 6 & 8 \\ \hline \end{array} \end{array} \xrightarrow{R_3 - R_1} \begin{array}{c} \underline{\underline{B}} \\ \begin{array}{|ccc|c} \hline 1 & 2 & 3 & 4 \\ \hline 0 & 1 & 2 & 3 \\ \hline 0 & 3 & 3 & 4 \\ \hline \end{array} \end{array}$$

We can write this operation also as

$$\begin{array}{|c|} \hline \underline{\underline{A}} \\ \hline \end{array} \begin{array}{l} R_1 \rightarrow 1 \cdot R_1 + 0 \cdot R_2 + 0 \cdot R_3 \\ R_2 \rightarrow 0 \cdot R_1 + 1 \cdot R_2 + 0 \cdot R_3 \\ R_3 \rightarrow -1 \cdot R_1 + 0 \cdot R_2 + 1 \cdot R_3 \end{array} \begin{array}{|c|} \hline \underline{\underline{B}} \\ \hline \end{array}$$
$$\begin{array}{|ccc|} \hline 1 & 0 & 0 \\ \hline 0 & 1 & 0 \\ \hline -1 & 0 & 1 \\ \hline \end{array} = \underline{\underline{C}}$$

The coefficients form a matrix themselves and more over

$$\underline{\underline{C}} \cdot \underline{\underline{A}} = \underline{\underline{B}}.$$

Every row operation we've performed is equivalent to a left multiplication by an "invertible"  $n \times n$  matrix.

The process of elimination of variables from the viewpoint of matrices is the following.

(1) You write your system in matrix form as follows

$$\begin{cases} a_{11}x_1 + \dots + a_{1n}x_n = b_1 \\ a_{21}x_1 + \dots + a_{2n}x_n = b_2 \\ \vdots \\ a_{m1}x_1 + \dots + a_{mn}x_n = b_n \end{cases} \iff \underline{\underline{A}} \cdot \underline{\underline{x}} = \underline{\underline{b}}$$

$\downarrow$   
where  $[\underline{\underline{A}}]_{ij} = a_{ij}$

(2) Multiply both sides of the equation with well-chosen matrices  $\underline{\underline{C}}_1, \underline{\underline{C}}_2, \underline{\underline{C}}_3, \dots, \underline{\underline{C}}_k$  that perform the row operations. Let  $\underline{\underline{C}} = \underline{\underline{C}}_k \cdot \underline{\underline{C}}_{k-1} \cdot \dots \cdot \underline{\underline{C}}_2 \cdot \underline{\underline{C}}_1$  then the new system becomes <sup>to bring the LHS in RREF.</sup>

$$(\underline{\underline{C}} \cdot \underline{\underline{A}}) \underline{\underline{x}} = \underline{\underline{C}} \cdot \underline{\underline{b}}$$

(3) Once  $(\underline{\underline{C}} \cdot \underline{\underline{A}})$  is in RREF we convert back to a system and set up the solutions.

SPECIAL SCENARIO If  $\underline{\underline{A}}$  is an  $n \times n$  matrix and there exists a sequence of row operations to transform the system

$$\left[ \begin{array}{c|c} \underline{\underline{A}} & \underline{\underline{b}} \end{array} \right]$$

→

Unit matrix  
 $\underline{\underline{I}}_{n \times n}$

$$\left[ \begin{array}{c|c} \begin{matrix} 1 & 0 & \dots & 0 \\ 0 & 1 & & \vdots \\ \vdots & & \ddots & 0 \\ 0 & \dots & & 1 \end{matrix} & \begin{matrix} \sim \underline{\underline{b}} \\ \sim \underline{\underline{b}} \\ \vdots \\ \sim \underline{\underline{b}} \end{matrix} \end{array} \right] = \underline{\underline{C}} \cdot \underline{\underline{b}} = \underline{\underline{A}}^{-1} \cdot \underline{\underline{b}}$$

then  $\underline{\underline{A}}$  is an invertible matrix and the matrix

$\underline{\underline{C}} = \underline{\underline{C}}_k \cdot \underline{\underline{C}}_{k-1} \cdot \dots \cdot \underline{\underline{C}}_2 \cdot \underline{\underline{C}}_1$  used to bring the system into RREF is the inverse of  $\underline{\underline{A}}$ ;  $\underline{\underline{A}}^{-1}$ .

With other words, if the system of equations can be written in matrix form like

$$\underline{A} \underline{x} = \underline{b}$$

where  $\underline{A}$  is an  $n \times n$  invertible matrix then we can solve the system by multiplying on the left by  $\underline{A}^{-1}$  to get

$$\underline{A}^{-1} \underline{A} \underline{x} = \underline{A}^{-1} \underline{b}$$

$$\Leftrightarrow \underline{I}_{n \times n} \underline{x} = \underline{A}^{-1} \underline{b}$$

$$\underline{x} = \underline{A}^{-1} \underline{b}$$

**EXAMPLE:** Given that  $\begin{bmatrix} 9 & -4 \\ -2 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 4 \\ 2 & 9 \end{bmatrix}$  solve  $\begin{cases} 9x - 4y = 2 \\ -2x + y = -1 \end{cases}$

**Q:** how to find  $\underline{A}^{-1}$ ? (if it exists)

**A:** When we reduce  $\underline{A}$  to RREF we can keep track of the matrix used to transform  $\underline{A}$  to  $\underline{I}_{n \times n}$  as follows. Rather than row reducing  $\underline{A}$  we row reduce the matrix

$\underline{A}$	$\underline{I}_{n \times n}$
-----------------	------------------------------

Row reducing this extended matrix is the same as multiplying both matrices simultaneously by the same matrices  $\underline{C}_1, \underline{C}_2, \dots, \underline{C}_k$ , so we get a new matrix that looks like

$$\underline{C} \underline{A} = \underline{I}_{n \times n} \quad \text{so} \quad \underline{C} = \underline{A}^{-1}$$

$$\begin{array}{c} \downarrow \\ \begin{array}{|c|c|} \hline \underline{C} \underline{A} & \underline{C} \underline{I}_{n \times n} \\ \hline \end{array} = \begin{array}{|c|c|} \hline \underline{I}_{n \times n} & \underline{C} \\ \hline \end{array} \end{array}$$

# EXAMPLES (1) Solve the system of lin eqns

$$\begin{cases} 5x + 4y = 1 \\ 6x + 5y = 2 \end{cases}$$

by (a) rewriting it in the form

$$\underline{\underline{A}} \underline{\underline{x}} = \underline{\underline{b}}, \text{ (b) finding } \underline{\underline{A}}^{-1}, \text{ and } \textcircled{c} \text{ finding } \underline{\underline{x}} \text{ via } \underline{\underline{x}} = \underline{\underline{A}}^{-1} \underline{\underline{b}}$$

$\downarrow$   
 $\begin{bmatrix} x \\ y \end{bmatrix}$

(a)  $\begin{bmatrix} 5 & 4 \\ 6 & 5 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$

(b) to find  $\underline{\underline{A}}^{-1}$ , let's bring  $\begin{bmatrix} 5 & 4 & 1 & 0 \\ 6 & 5 & 0 & 1 \end{bmatrix}$  in RREF.

$$\begin{bmatrix} 5 & 4 & 1 & 0 \\ 6 & 5 & 0 & 1 \end{bmatrix} \xrightarrow{\substack{R_1 \cdot 6 \\ R_2 \cdot 5}} \begin{bmatrix} 30 & 24 & 6 & 0 \\ 30 & 25 & 0 & 5 \end{bmatrix}$$

$$\xrightarrow{R_2 - R_1} \begin{bmatrix} 30 & 24 & 6 & 0 \\ 0 & 1 & -6 & 5 \end{bmatrix} \xrightarrow{R_1 - 24 \cdot R_2} \begin{bmatrix} 30 & 0 & 25 \cdot 6 & -24 \cdot 5 \\ 0 & 1 & -6 & 5 \end{bmatrix}$$

$$\xrightarrow{R_1 / 30} \begin{bmatrix} 1 & 0 & 5 & -4 \\ 0 & 1 & -6 & 5 \end{bmatrix} \text{ so } \underline{\underline{A}}^{-1} = \begin{bmatrix} 5 & -4 \\ -6 & 5 \end{bmatrix}$$

(c)  $\begin{bmatrix} x \\ y \end{bmatrix} = \underline{\underline{x}} = \underline{\underline{A}}^{-1} \underline{\underline{b}} = \begin{bmatrix} 5 & -4 \\ -6 & 5 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} -3 \\ 4 \end{bmatrix}$

(2) Find the inverse of  $\underline{\underline{A}} = \begin{bmatrix} -2 & 3 & 1 \\ 1 & -1 & -2 \\ -1 & 3 & -7 \end{bmatrix}$

$$\begin{bmatrix} -2 & 3 & 1 & 1 & 0 & 0 \\ 1 & -1 & -2 & 0 & 1 & 0 \\ -1 & 3 & -7 & 0 & 0 & 1 \end{bmatrix} \xrightarrow{R_1 \leftrightarrow R_2} \begin{bmatrix} 1 & -1 & -2 & 0 & 1 & 0 \\ -2 & 3 & 1 & 1 & 0 & 0 \\ -1 & 3 & -7 & 0 & 0 & 1 \end{bmatrix}$$

$$\begin{array}{l} R_2 + 2R_1 \\ R_3 + R_1 \end{array} \rightarrow \begin{array}{|cccccc|} \hline 1 & -1 & -2 & 0 & 1 & 0 \\ \hline 0 & 1 & -3 & 1 & 2 & 0 \\ \hline 0 & 2 & -9 & 0 & 1 & 1 \\ \hline \end{array}$$

$$\begin{array}{l} R_3 - 2R_2 \end{array} \rightarrow \begin{array}{|cccccc|} \hline 1 & -1 & -2 & 0 & 1 & 0 \\ \hline 0 & 1 & -3 & 1 & 2 & 0 \\ \hline 0 & 0 & -3 & -2 & -3 & 1 \\ \hline \end{array} \xrightarrow{R_3 \cdot \left(\frac{1}{3}\right)} \begin{array}{|cccccc|} \hline 1 & -1 & -2 & 0 & 1 & 0 \\ \hline 0 & 1 & -3 & 1 & 2 & 0 \\ \hline 0 & 0 & 1 & \frac{2}{3} & 1 & -\frac{1}{3} \\ \hline \end{array}$$

$$\begin{array}{l} R_1 + 2R_2 \\ R_2 + 3R_3 \end{array} \rightarrow \begin{array}{|cccccc|} \hline 1 & -1 & 0 & \frac{4}{3} & 3 & -\frac{2}{3} \\ \hline 0 & 1 & 0 & 3 & 5 & -1 \\ \hline 0 & 0 & 1 & \frac{2}{3} & 1 & -\frac{1}{3} \\ \hline \end{array}$$

$$R_1 + R_2 \rightarrow \begin{array}{|ccc|ccc|} \hline 1 & 0 & 0 & \frac{13}{3} & 8 & -\frac{5}{3} \\ \hline 0 & 1 & 0 & 3 & 5 & -1 \\ \hline 0 & 0 & 1 & \frac{2}{3} & 1 & -\frac{1}{3} \\ \hline \end{array}$$

$$\text{So } \underline{A}^{-1} = \begin{array}{|ccc|} \hline \frac{13}{3} & 8 & -\frac{5}{3} \\ \hline 3 & 5 & -1 \\ \hline \frac{2}{3} & 1 & -\frac{1}{3} \\ \hline \end{array}$$

## SOLVING MULTIPLE SYSTEMS AT ONCE

If you have multiple systems of lin equations whose coefficient matrix is the same matrix  $\underline{A}$ , so

$$\underline{A} \underline{x}_1 = \underline{b}_1, \quad \underline{A} \underline{x}_2 = \underline{b}_2, \quad \dots, \quad \underline{A} \underline{x}_k = \underline{b}_k$$

then you can write this system as 1 matrix equation as follows:

$$\underline{A} \cdot \underbrace{\begin{array}{|c|c|c|} \hline \underline{x}_1 & \underline{x}_2 & \dots & \underline{x}_k \\ \hline \end{array}}_{\underline{X}} = \underbrace{\begin{array}{|c|c|c|} \hline \underline{b}_1 & \underline{b}_2 & \dots & \underline{b}_k \\ \hline \end{array}}_{\underline{B}}$$

or

$$\underline{\underline{A}} \cdot \underline{\underline{x}} = \underline{\underline{B}}$$

⊥